

Abortion and Substance Abuse

- Substance abuse involves the misuse of substances (alcohol, tobacco, illegal and legal drugs, and/or other mood-altering substances) and it is implicated in numerous mental and physical health problems as well as countless contemporary social ills [1].
- Peer-reviewed research published in leading journals in the U.S. and internationally has shown a consistent and statistically significant association between abortion and substance abuse of various forms [1-20]. The increased risk of substance abuse is estimated to be as high as 6 times greater among women who have aborted compared to women who have not aborted. Several of the studies comprising the world literature used pregnant women who delivered as the comparison group, with a few studies employing unintended pregnancy carried to term as the comparison group.
- A new particularly disturbing wave of research has shown associations between abortion and increased risk of substance abuse during later pregnancies [1,6,7, 11-13, 15, 16].
- The choice to abort is frequently filled with conflicting emotions and external pressures, often with the resulting decision not reflective of the woman's true desire [1]. Women who are relatively comfortable with an abortion at the time of the procedure may come to deeply regret their choice in the months and years that follow. Under such circumstances, women may experience abortion as a significant loss involving grief and/or traumatic stress, guilt, anger, anxiety, depression, and other psychological symptoms that can precipitate substance abuse [1,21]. Dysfunctional coping mechanisms including substance abuse may be relied upon to self-medicate and/or numb the emotional pain.
- Untreated substance abuse may facilitate avoidance and hinder coming to terms with the abortion [1]. When the emotional pain of an abortion is denied, the many physical, psychological, relational, and practical problems associated with substance abuse typically introduce new sources of pain and suffering that may compound and exacerbate the trauma of abortion. If women are encouraged to explore ambivalent feelings and make choices that are consonant with their fundamental desires and values prior to an abortion, much of this cycle could be averted and women's health thereby enhanced.
- Professionals who work with women suffering from substance abuse disorders should sensitively inquire about a past abortion in order to provide women with an opportunity to express their feelings in a safe, comfortable context. Efforts directed toward treating women for substance use disorders without addressing what in many instances is a primary cause are inclined to be less effective and in some cases may prove futile[1].



1. Coleman, PK. Induced Abortion and Increased Risk of Substance Abuse: A Review of the Evidence. *Current Women's Health Reviews*. 2005;1:21-34.
2. Amaro H, Zuckerman B, Cabral H. Drug use among adolescent mothers: profile of risk. *Pediatrics* 1989;84:144-51.
3. Coleman PK. Resolution of Unwanted Pregnancy During Adolescence Through Abortion versus Childbirth: Individual and Family Predictors and Consequences. *Journal of Youth and Adolescence* 2006; 35, 903-911.
4. Coleman, PK, Coyle, CT, Shuping M, & Rue V. Induced Abortion and Anxiety, Mood, and Substance Abuse Disorders: Isolating the Effects of Abortion in the National Comorbidity Survey. *Journal of Psychiatric Research* 2009; 43 770-776.
5. Coleman PK, & Maxey DC, Spence M, & Nixon C. (in press). The choice to abort among mothers living under ecologically deprived conditions: Predictors and consequences. *International Journal of Mental Health and Addiction*.
6. Coleman PK, Reardon DC, Cogle J. Substance use among pregnant women in the context of previous reproductive loss and desire for current pregnancy. *British Journal of Health Psychology* 2005; 10 (2): 255-268 (14).
7. Coleman PK, Reardon DC, Rue V, Cogle J. Prior history of induced abortion in relation to substance use during subsequent pregnancies carried to term. *American Journal of Obstetrics and Gynecology* 2002;187:1673-78.
8. Dingle K, Alta R, Clavarino A et al. Pregnancy loss and psychiatric disorders in young women: An Australian birth cohort study. *The British Journal of Psychiatry* 2008; 193: 455-460.
9. Fergusson DM, Horwood JH, & Boden JM. Abortion and mental health disorders: Evidence from a 30-year longitudinal study, *The British Journal of Psychiatry* 2008; 193: 444-451.
10. Fergusson DM, Horwood J, & Ridder EM. Abortion in young women and subsequent mental health. *Journal of Child Psychology and Psychiatry* 2006; 47: 16-24.
11. Frank DA, Zuckerman BS, Amaro H, et al. Cocaine use during pregnancy: Prevalence and correlates. *Pediatrics* 1988; 82: 888-895.
12. Gladstone J, Levy M, Nulman I, et al. Characteristics of pregnant women who engage in binge alcohol consumption. *CMAJ* 1997;156:789-794.
13. Graham K, Koren G. Characteristics of pregnant women exposed to cocaine in Toronto between 1985 and 1990. *CMAJ* 1991; 144:563.
14. Hope TL, Wilder EI, & Watt, TT. The relationships among adolescent pregnancy, pregnancy resolution, and juvenile delinquency. *The Sociological Quarterly* 2003; 44: 555-576.
15. Kuzma J, Kissinger D. Patterns of alcohol and cigarette use in pregnancy. *Neurobehav Toxicol Teratol* 1981;3:211-21.
16. Oro AS, Dixon, SD. Prenatal cocaine and methamphetamine exposure: maternal and neo-natal correlates. *Pediatrics* 1987;111:571-578.
17. Pedersen W. Childbirth, abortion and subsequent substance use in young women: a population-based longitudinal study. *Addiction* 2008; 102 (12): 1971-78.
18. Reardon DC, Coleman PK, Cogle J. Substance use associated with prior history of abortion and unintended birth: A national cross sectional cohort study. *American Journal of Drug and Alcohol Abuse*. 2004;26:369-383.
19. Reardon DC, Ney P. Abortion and subsequent substance abuse. *American Journal of Drug & Alcohol Abuse*. 2000; 26: 61-75.
20. Yamaguchi D, Kandel D. Drug use and other determinants of premarital pregnancy and its outcome: A dynamic analysis of competing life events. *Journal of Marriage and the Family*
21. Rue V, Coleman P, Rue J, Reardon D. Induced abortion and traumatic stress: A preliminary comparison of American and Russian women. *Medical Science Monitor* 2004; 10: SR5-16.